

JOURNAL OF ALGEBRA 90, 372-374 (1984)

A Formula for the Separability Idempotent in the Tensor Square of a Field

A. LEBOW AND M. SCHREIBER

*The Rockefeller University, New York, New York 10021**Communicated by Walter Feit*

Received February 16, 1983

1. INTRODUCTION

We shall establish here an explicit formula for the separability idempotent $([1, 2])$ in a ring of the form $E \otimes_k E$, where E is a separable finite extension of a ground field k . The formula is as follows.

Let \mathbf{a} be a primitive element for E over k , and let $\mathbf{a}_1 = \mathbf{a}, \mathbf{a}_2, \mathbf{a}_3, \dots, \mathbf{a}_d$ be its conjugates in the splitting field Ω of the minimal polynomial of \mathbf{a} . Let $V(\mathbf{a})$ be the matrix

$$V(\mathbf{a}) = \begin{pmatrix} 1 & \mathbf{a}_1 & \mathbf{a}_1^2 & \mathbf{a}_1^3 & \cdots & \mathbf{a}_1^{d-1} \\ 1 & \mathbf{a}_2 & \mathbf{a}_2^2 & \mathbf{a}_2^3 & \cdots & \mathbf{a}_2^{d-1} \\ & & *** & & & \\ 1 & \mathbf{a}_d & \mathbf{a}_d^2 & \mathbf{a}_d^3 & \cdots & \mathbf{a}_d^{d-1} \end{pmatrix}.$$

This defines a linear operator in Ω^d which is invertible by the assumed separability. The entries in ${}^tV(\mathbf{a}) \cdot V(\mathbf{a})$ are the power sums of the conjugates and so lie in k . Therefore the tensor θ defined as

$$\theta = \sum \{ {}^tV(\mathbf{a}) V(\mathbf{a}) \}^{-1} {}_{ij}(\mathbf{a}^i \otimes \mathbf{a}^j) \quad (1)$$

is an element of $E \otimes_k E$. We will prove that it is the separability idempotent in $E \otimes_k E$. We will also show how the use of a primitive element may be avoided at the expense of further calculation.

2. PRELIMINARIES

We write $\mathbf{G}(K/F)$ for the group of automorphisms of a field $K \supset F$ fixing F pointwise, and id for the identity map. Any two maps $f, g: K \rightarrow K$ fixing F determine a map $f \times g: K \otimes_F K \rightarrow K$ defined by the relation $f \times g(x \otimes y) = f(x)g(y)$.

We recall that if E is a separable finite extension of k then $E \otimes_k E$ is the algebra-direct-sum $E \otimes_k E = \bigoplus_j (E \otimes_k E) e_j$ of its minimal ideals $(E \otimes_k E) e_j$, each principal, generated by a uniquely determined indecomposable idempotent e_j ; that these idempotents satisfy $e_j e_k = \delta_{jk} e_k$ and $\sum e_j = 1 \otimes 1$; that each minimal ideal is a field; and that all ideals in $E \otimes_k E$ have unique complements.

By the *support ideal* of a ring homomorphism $h: E \otimes_k E \rightarrow M$, M any extension of E , we mean the ideal complementary to $\ker(h)$. We say $T \in E \otimes_k E$ satisfies the *symmetry condition* if

$$(1 \otimes x - x \otimes 1)T = 0, \quad \text{all } x \in E. \quad (2)$$

For any separable finite extension E of k there is precisely one idempotent in $E \otimes_k E$ satisfying (2), namely, the generator of the support ideal of $\text{id} \times \text{id}$. That idempotent is the *separability idempotent* in $E \otimes_k E$. It is indecomposable because the support ideal of $\text{id} \times \text{id}$ is a field.

3. PROOF OF THE FORMULA

With matters and notation as in Section 1, we interpret θ as an element of $\Omega \otimes_k \Omega$, and assert that, as such,

$$(\text{id} \times \sigma)(\theta) = \begin{cases} 1 & \text{if } \sigma \in \mathbf{G}(\Omega/E) \\ 0 & \text{if } \sigma \notin \mathbf{G}(\Omega/E). \end{cases} \quad (3)$$

To prove (3) we put

$$\bar{\mathbf{a}}_k = (1, \mathbf{a}_k, \mathbf{a}_k^2, \dots, \mathbf{a}_k^{d-1}).$$

Then, if $\{\bar{\mathbf{b}}_k\}_1^d$ is that basis of Ω^d such that $(\bar{\mathbf{b}}_k)_j = \delta_{kj}$, we have

$${}^t V(\mathbf{a}) \bar{\mathbf{b}}_k = \bar{\mathbf{a}}_k.$$

For $\sigma \in \mathbf{G}(\Omega/k)$ we have

$$\begin{aligned} (\text{id} \times \sigma)(\theta) &= \sum_{ij} \{ {}^t V(\mathbf{a}) V(\mathbf{a}) \}^{-1}_{ij} (\mathbf{a}_1^i \cdot \sigma(\mathbf{a}_1^j)) \\ &= \langle {}^t V(\mathbf{a})^{-1} {}^t V(\mathbf{a})^{-1} \bar{\mathbf{a}}_1, \sigma(\bar{\mathbf{a}}_1) \rangle \\ &= \langle {}^t V(\mathbf{a})^{-1} \bar{\mathbf{a}}_1, {}^t V(\mathbf{a})^{-1} \sigma(\bar{\mathbf{a}}_1) \rangle \end{aligned}$$

in the natural scalar product in Ω^d . Now $\sigma(\bar{\mathbf{a}}_1) = \bar{\mathbf{a}}_1$ if $\sigma \in \mathbf{G}(\Omega/E)$, and $\sigma(\bar{\mathbf{a}}_1) = \bar{\mathbf{a}}_k$, $k \neq 1$, if $\sigma \notin \mathbf{G}(\Omega/E)$ because $E = k(\mathbf{a}_1)$. In the first case $(\text{id} \times \sigma)(\theta) = \langle {}^t V(\mathbf{a})^{-1} \bar{\mathbf{a}}_1, {}^t V(\mathbf{a})^{-1} \bar{\mathbf{a}}_1 \rangle = \langle \bar{\mathbf{b}}_1, \bar{\mathbf{b}}_1 \rangle = 1$. In the second case $(\text{id} \times \sigma)(\theta) = \langle \bar{\mathbf{b}}_1, \bar{\mathbf{b}}_k \rangle = 0$. This completes the proof of (3).

Since Ω is a Galois extension of k the set of indecomposable idempotents in $\Omega \otimes_k \Omega$ is the set $\{e_\sigma: \sigma \in \mathbf{G}(\Omega/k)\}$ where $e_\sigma = (\sigma \otimes \text{id})e_1$, e_1 being the separability idempotent in $\Omega \otimes_k \Omega$. Thus $\Omega \otimes_k \Omega = \bigoplus_\sigma (\Omega \otimes_k \Omega)e_\sigma$, and it is immediate (from the fact that $(1 \otimes x - x \otimes 1)e_1 = 0$ for all $x \in \Omega$) that $(\Omega \otimes_k \Omega)e_\sigma = (\Omega \otimes_k 1)e_\sigma$. Reasoning from the way in which $\mathbf{G}(\Omega/k)$ acts upon $\{e_\sigma\}$, namely, $(\theta \otimes \eta)e_\sigma = e_{(\theta\sigma\eta^{-1})}$, one finds that the support ideal of $\text{id} \times \sigma$ is $(\Omega \otimes_k \Omega)e_\sigma$. Writing $T \in \Omega \otimes_k \Omega$ as $T = \sum (T_\tau \otimes 1)e_\tau$, $T_\tau \in \Omega$, and applying $\text{id} \times \sigma$ to both sides, we then have $T_\sigma = (\text{id} \times \sigma)T$. In these terms (3) is the statement that $\Theta = \sum \{e_\sigma: \sigma \in \mathbf{G}(\Omega/E)\}$. In particular we thus know Θ to be idempotent. It remains only to check that Θ satisfies the symmetry condition (2) in $E \otimes_k E$. We do this by viewing $T = (1 \otimes x - x \otimes 1)\Theta$, $x \in E$, as an element of $\Omega \otimes_k \Omega$ and showing that all its "coordinates" $(\text{id} \times \sigma)T = T_\sigma$ vanish. We have $(\text{id} \times \sigma)T = (\sigma(x) - x) \cdot (\text{id} \times \sigma)(\Theta)$. Now the first factor vanishes if $\sigma \in \mathbf{G}(\Omega/E)$, and by (3) the second factor vanishes if $\sigma \notin \mathbf{G}(\Omega/E)$. Therefore the product vanishes. This completes the proof.

We remark in closing that the use of a primitive element in the construction of the separability idempotent may be avoided as follows. If E is constructed by a series of adjunctions, say $E = k(r_1, r_2, \dots, r_n)$, then the separability idempotent R_j in each of the intermediate algebras $k(r_j) \otimes_k k(r_j)$ is given by (1) with $\mathbf{a} = r_j$, and the product in $E \otimes_k E$ of these idempotents is the separability idempotent in $E \otimes_k E$. To see this it is sufficient to check that

$$(1 \otimes r_i - r_i \otimes 1) \left(\prod_j R_j \right) = 0, \quad \text{all } i.$$

But $E \otimes_k E$ is commutative, and $(1 \otimes r_i - r_i \otimes 1)R_i = 0$ for all i , so $(1 \otimes r_i - r_i \otimes 1)(\prod_j R_j) = (\prod_{j \neq i} R_j)(1 \otimes r_i - r_i \otimes 1)R_i = 0$ for each i .

ACKNOWLEDGMENT

We thank H. Bass, D. Blasius, and S. Kochen for helpful conversations, and we thank the referee for catching an error in the first version of this paper.

REFERENCES

1. F. DEMEYER AND E. INGRAHAM, "Separable Algebras over Commutative Rings," Lecture Notes in Mathematics No. 181, Springer-Verlag, New York, 1971.
2. S. U. CHASE, D. K. HARRISON, AND A. ROSENBERG, Galois theory and cohomology of commutative rings, *Mem. Amer. Math. Soc.* **52** (1965).